
Drug Checking in British Columbia


of Samples Tested by Expected Drug Category:

265 Samples Tested

Opioid: 145 / Stimulant: 48 / Psychedelic: 57 / Unknown: 11 / Depressant: 3 / Other: 0 / Polysubstance: 1







Public health notifications Expected Fentanyl Date & Location **Drugs Detected** What does this mean? Drug Strip 8-August-2019, SafePoint, Surrey Down N/A N/A Increased risk of overdoses; don't use alone and carry naloxone 14-August-2019, Insite, Vancouver Down Etizolam Positive Benzos may increase the risk of an overdose; use at an OPS or SCS Crack 24-August-2019, SafePoint, Surrey Heroin and Fentanyl Positive Check your drugs for fentanyl at any OPS or SCS Cocaine To receive drug contamination alerts in the VCH region, text "alert" to (236) 999-DOPE (3673)

This report summarizes key indicators for Fourier Transform Infrared (FT-IR)/fentanyl test strip combination drug checking done in British Columbia in August 2019.

As the FT-IR spectrometer has a detection limit of 3-4% by weight (Sherman et al., 2018), unless otherwise stated, a drug check on any given sample consists of <u>both</u> Bruker ALPHA Fourier-Transform Infrared (FTIR) spectrometer <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine or lysergic acid diethylamide (LSD) immunoassay test strips are also used.

Please note that these data are not finalized and are subject to change.

^{* &}quot;Fentanyl not tested" means that a fentanyl test stip wasn't done for the sample because the client declined to do the test, or they left before the test was able to be completed.

Drug Checking in British Columbia

of Samples Tested by Region:

Total #: 265

216: Vancouver Coastal Health region (82%)



9: Interior Health region (3%)

40: Fraser Health region (15%)

of samples that matched client expectation

*using FTIR/test strip drug checking

EXPECTED DRUG: Depressant 3 Samples Tested



Matched: 1 Did not match: 2





Match not

EXPECTED DRUG: Psychedelic 57 Samples Tested

EXPECTED DRUG:

145 Samples Tested

Opioid



Matched: 54 Did not match: 3

Matched: 118 Did not match: 27 EXPECTED DRUG: **Stimulant** 48 Samples Tested

> Matched: 47 Did not match: 1

EXPECTED DRUG: **Polysubstance**


1 Sample Tested



Matched: 1


determined: 11

Total 265 Samples Tested



Matched: 221 Did not match: 33 Match not determined: 11

Drug Checking in British Columbia



Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

*'Down' can refer to any opioid drug present in any amount

Depressant may include:

benzodiazepines, GHB, zopiclone **Opioid may include:** "down," heroin, fentanyl, fentanyl analogues, pharmaceutical opioids

Polysubstance includes: cross-category mixtures

Psychedelic may include: 2C-family, tryptamines, ketamine, MDMA, LSD, cannabis Stimulant may include: "speed", crystal meth, cocaine, cathinones Unknown includes: all samples in which the client could not identify the expected substance

BCCSU gratefully acknowledges the contributions of the following partners:

Portland Hotel Society | Insite | Molson Overdose Prevention Site | SisterSpace | Overdose Prevention Society | SafePoint
The Abbotstford Community Hub | ANKORS | ASK Wellness | St. Paul's Overdose Prevention Site | Powell Street Getaway | Three Bridges



























