
## **Drug Checking in British Columbia**

### # of Samples Tested by Expected Drug Category:

377 Samples Tested

Opioid: 224 / Stimulant: 57 / Psychedelic: 64 / Unknown: 21 / Depressant: 6 / Other: 4 / Polysubstance: 1





#### # of samples tested for fentanyl **64** Psychedelic **57** Stimulant 224 Opioid 6 Depressant 1 Polysubstance 21 Unknown 4 Other samples samples samples samples samples sample samples **55** FENTANYL NEGATIVE 193 FENTANYL POSITIVE 52 FENTANYL NEGATIVE 18 FENTANYL NEGATIVE 3 FENTANYL NEGATIVE (81%)(96%)(86%)28 **FENTANYL** FENTANYL 12 FENTANYL NOT FENTANYL NOT FENTANYI NOT 1 FENTANYL POSITIVE 3 FENTANYL POSITIVE 6 FENTANYI NEGATIVE 1 FENTANYL NOT TESTED\* POSITIVE TESTED\* **NEGATIVE** TESTED\* TESTED\* (100%)(100%) (14%) (25%) (13%) (19%) (1%) (2%)



### **Public Health Notifications**

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	What does this mean?
12 September 2019- Insite	Fentanyl	Suspected Carfentanil, Etizolam	Positive	May cause prolonged ODs; Check your drugs at any OPS or SCS

#### To receive drug contamination alerts in the VCH region, text "alert" to (236) 999-DOPE (3673)

This report summarizes key indicators for Fourier Transform Infrared (FT-IR)/fentanyl test strip combination drug checking done in British Columbia in September 2019.

As the FT-IR spectrometer has a detection limit of 3-4% by weight (Sherman et al., 2018), unless otherwise stated, a drug check on any given sample consists of <u>both</u> Bruker ALPHA Fourier-Transform Infrared (FTIR) spectrometer <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine or lysergic acid diethylamide (LSD) immunoassay test strips are also used.

Please note that these data are not finalized and are subject to change.

<sup>\* &</sup>quot;Fentanyl not tested" means that a fentanyl test stip wasn't done for the sample because the client declined to do the test, or they left before the test was able to be completed.

## **Drug Checking in British Columbia**

#### # of Samples Tested by Region:

Total #: 377

249: Vancouver Coastal Health region (66%)



70: Interior Health region (19%)

58: Fraser Health region (15%)

## # of samples that matched client expectation

\*using FTIR/test strip drug checking

**EXPECTED DRUG: Depressant** 

6 Samples Tested



Matched: 3 Did not match: 3

**EXPECTED DRUG:** Unknown 21 Samples Tested



Match not determined: 21

**EXPECTED DRUG: Psychedelic** 

64 Samples Tested



Matched: 49 Did not match: 11 Match not determined: 4 EXPECTED DRUG: **Polysubstance** 

1 Sample Tested



Matched: 1

**EXPECTED DRUG: Stimulant** 



Matched: 55 Did not match: 2

EXPECTED DRUG:

**Opioid** 

224 Samples Tested



Matched: 205 Did not match: 18 Match not determined: 1 EXPECTED DRUG: Other

4 Samples Tested



Matched: 4

**Total** 


377 Samples Tested



Matched: 317 Did not match: 34

Match not determined: 26

# **Drug Checking in British Columbia**



Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents
\*'Down' can refer to any opioid drug present in any amount

Depressant may include:

benzodiazepines, GHB, zopiclone Opioid may include: "down," heroin, fentanyl, fentanyl analogues, pharmaceutical opioids Polysubstance includes: cross-category mixtures

Psychedelic may include: 2C-family, tryptamines, ketamine, MDMA, LSD, cannabis Stimulant may include: "speed", crystal meth, cocaine, cathinones Unknown includes: all samples in which the client could not identify the expected substance

#### BCCSU gratefully acknowledges the contributions of the following partners:

Portland Hotel Society | Insite | Molson Overdose Prevention Site | SisterSpace | Overdose Prevention Society | SafePoint
The Abbotstford Community Hub | ANKORS | ASK Wellness | St. Paul's Overdose Prevention Site | Powell Street Getaway | Three Bridges



























