SAMPLE PRESCRIBER'S ORDERS

This is an example of a pre-printed order that may be used in the provision of iOAT.

Patient label	
---------------	--

Date and	TITRATION ORDERS FOR DIACETYLMORPHINE
time	(Items with checkboxes must be selected to be ordered)
	These orders are applicable to clients who are starting on diacetylmorphine treatment.
	INDICATION: Opioid use disorder
	MEDICATION: Diacetylmorphine 100mg/mL injection for the following orders for ALL routes of administration (IM/IV)
	Qualified health professional to assess for signs and symptoms of opioid sedation and withdrawal before and after each dose using Pasero Opioid-induced Sedation Scale (POSS) If POSS 3 or 4, notify prescriber immediately and hold next dose
	 □ Qualified health professional to assess for signs and symptoms of stimulant use before each dose and notify prescriber if the following symptoms are noted: □ Qualified health professional to assess for alcohol use with breathalyzer before each dose. Hold dose and notify prescriber if breathalyzer reading is: ■ Rapid urine drug testing PRN ■ Breathalyzer PRN
	 Breathalyzer PRN Qualified health professional to notify prescriber if client misses 2 or more sessions or declines second dose
	 Qualified health professional to notify prescriber about titration by 3pm on Day 2 in order for prescriber to write maintenance prescription Round all doses UP to nearest 5mg
	Day 1, Session 1: Diacetylmorphine 15mg (fifteen milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose
	Dose 2: Diacetylmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose Qualified health professional to advise client of timing for return for Session 2
	Do not exceed 45mg (forty-five milligrams) of diacetylmorphine IM/IV for Day 1, Session 1
	Day 1, Session 2: Dose 1: Diacetylmorphine 45mg (forty-five milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose

THIS IS A SAMPLE PRE-PRINTED ORDER ADAPTED FROM CROSSTOWN CLINIC IN VANCOUVER, BC.
THIS FORM IS NOT MEANT FOR CLINICAL USE.





SAMPLE PRESCRIBER'S ORDERS

This is an example of a pre-printed order that may be used in the provision of iOAT.

Dose 2: Diacetylmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose Qualified health professional to advise client of timing for return for Session 3

Do not exceed 75mg (seventy-five milligrams) of diacetylmorphine IM/IV for Day 1, Session 2

Day 1, Session 3:

Dose 1: Diacetylmorphine 75mg (seventy-five milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: Diacetylmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Day 2

Do not exceed 105mg (one hundred and five milligrams) of diacetylmorphine IM/IV for Day 1, Session 3

Day 2, Session 1:

Dose 1: Diacetylmorphine 40% of total daily dose from DAY 1 to a maximum of 90mg (ninety milligrams)

IV/IM for client self-administration x 1 dose

Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: Diacetylmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose Qualified health professional to advise client of timing for return for Session 2

Do not exceed 120mg (one hundred and twenty milligrams) of diacetylmorphine IM/IV for Day 2, Session 1

Day 2, Session 2:

Dose 1: If earlier doses were well tolerated:

Diacetylmorphine maximum tolerated Day 2, Session 1 dose to a maximum of 120mg (one hundred and twenty milligrams) IV/IM for client self-administration x 1 dose

Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: Diacetylmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose Qualified health professional to advise client of timing for return for Session 3

Do not exceed 150mg (one hundred and fifty milligrams) of diacetylmorphine IM/IV for Day 2, Session 2

Day 2, Session 3:

Dose 1: If earlier doses were well tolerated:

Diacetylmorphine maximum tolerated Day 2, Session 2 dose to a maximum of 150mg (one hundred and fifty milligrams) IV/IM for client self-administration x 1 dose

THIS IS A SAMPLE PRE-PRINTED ORDER ADAPTED FROM CROSSTOWN CLINIC IN VANCOUVER, BC.
THIS FORM IS NOT MEANT FOR CLINICAL USE.





SAMPLE PRESCRIBER'S ORDERS

This is an example of a pre-printed order that may be used in the provision of iOAT.

Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: Diacetylmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose Qualified health professional to advise client of timing for return for Day 3

Do not exceed 180mg (one hundred and eighty milligrams) of diacetylmorphine IM/IV for Day 2, Session 3

Day 3, Session 1:

Dose 1: Diacetylmorphine maximum tolerated Day 2, Session 3 dose to a maximum of 180mg (one hundred and eighty milligrams) IV/IM for client self-administration x 1 dose

Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Qualified health professional to advise client of timing for return for Session 2

Do not exceed 180mg (one hundred and eighty milligrams) of diacetylmorphine IM/IV for Day 3, Session 1

Day 3, Session 2:

Dose 1: Diacetylmorphine maximum tolerated Day 2, Session 3 dose to a maximum of 180mg (one hundred and eighty milligrams) IV/IM for client self-administration x 1 dose

Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Qualified health professional to advise client of timing for return for Session 3

Do not exceed 180mg (one hundred and eighty milligrams) of diacetylmorphine IM/IV for Day 3, Session 2

Day 3, Session 3:

Dose 1: Diacetylmorphine maximum tolerated Day 2, Session 3 dose to a maximum of 180mg (one hundred and eighty milligrams) IV/IM for client self-administration x 1 dose

Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Qualified health professional to advise client of timing for return to clinic now that titration is complete

Do not exceed 180mg (one hundred and eighty milligrams) of diacetylmorphine IM/IV for Day 3, Session 3





