

# **Drug Checking in Vancouver Coastal Health**

### # of Samples Tested by Expected Drug Category:


535 Samples Tested \*


Psychedelic: 124 Unknown: 28 / Depressant: 18 / Other: 5 / Polysubstance: 1








## # of samples tested for fentanyl






















## **Public Health Notifications**

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	What does this mean?
26 February 2020- Insite, Vancouver	Down	Caffeine, sugars, Fentanyl or analogue	Positive	Positive	Benzos can cause prolonged ODs
26 February 2020- Insite, Vancouver	Down	Caffeine, Mannitol and inositol (sugars), Fentanyl HCl	Positive	Positive	Check for benzos in your drugs at any OPS/SCS
21 February 2020 - OPS, Vancouver	Down	Caffeine, Erythritol (sugar)	Positive	Positive	Test your drugs for fentanyl at any local OPS or SCS
14 February 2020- OPS, Vancouver	Down	Caffeine, Mannitol, Fentanyl or analogue	Positive	Positive	Use at an OPS or SCS
7 February 2020- OPS, Vancouver	Down	Caffeine, Mannitol, Fentanyl or analogue	Positive	Positive	Benzo-complicated ODs can recur after Naloxone
6 February 2020- OPS, Vancouver	Fentanyl	Caffeine, Mannitol, Fentanyl	Positive	Positive	Test your drugs!

6.1% of expected opioid samples tested positive for benzodiazepines in the Vancouver Coastal Region region

#### To receive drug contamination alerts, text "alert" to (236) 999-DOPE (3673)

This report summarizes key indicators for Fourier Transform Infrared (FTIR)/fentanyl test strip combination drug checking done in Vancouver Coastal Health in February 2020. As there is 5% fentanyl detection limit on the Bruker ALPHA Fourier-Transform Infrared (FTIR) spectrometer (McCrae, 2019), a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine or lysergic acid diethylamide (LSD) immunoassay test

Benzodiazepine-positivity of opioid samples are determined using BTNX benzodiazeoine test strips, which have not been formally evaluated for drug checking. Therefore, the results presented above are presumptive and tentative until confirmed by a laboratory.

\* "Fentanyl not tested" means that a fentanyl test stip wasn't done for the sample because the client declined to do the test, or they left before the test was able to be completed.

\* Please note that reports now include sample data collected at the Get Your Drugs Tested site located in Vancouver and that data are not finalized and are subject to



strips are also used.




## # of drugs checks and hours at each respective site:

Location	# of Drug Checks	Hours
Insite	20	6
Molson OPS	28	9
Overdose Prevention Society OPS	43	14
St. Paul's OPS	14	7
Dr. Peter Centre	11	4
Get Your Drugs Tested	419	232
Total	535	272


# # of samples that matched client expectation

\*using FTIR/test strip drug checking



#### **Total**

535 Samples Tested



# **Drug Checking in Vancouver Coastal Health**

### # of opioid samples that matched client expectation

\*using FTIR/test strip drug checking

EXPECTED DRUG: Heroin

9 Samples Tested




Matched: 8
Did not match: 1

EXPECTED DRUG:

Down

104 Samples Tested



Matched: 97
Did not match: 7

**EXPECTED DRUG:** 

**Fentanyl and Heroin** 

2 Samples Tested



Matched: 2

**EXPECTED DRUG:** 

**Pharmaceutical** 

14 Samples Tested



Matched: 5

Did not match: 2

Match not determined: 7

**EXPECTED DRUG:** 

**Fentanyl** 

137 Samples Tested



Matched: 124 Did not match: 13

**Total** 

266 Samples Tested



Matched: 236
Did not match: 23
Match not determined: 7

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

\*'Down' can refer to any opioid drug present in any amount

Depressant may include: benzodiazepines, GHB, zopiclone Opioid may include: "down," heroin, fentanyl, fentanyl analogues, pharmaceutical opioids Polysubstance includes: cross-category mixtures

Psychedelic may include: 2C-family, tryptamines, ketamine, MDMA, LSD, cannabis Stimulant may include: "speed", crystal meth, cocaine. cathinones Unknown includes: all samples in which the client could not identify the expected substance

**(ARMIK** 

BCCSU and VCH gratefully acknowledge the contributions of the following partners:

Portland Hotel Society | Insite | Molson Overdose Prevention Site | SisterSpace | Overdose Prevention Society

St. Paul's Overdose Prevention Site | Powell Street Getaway | Three Bridges







**%**Trovidence







